

Remedial Investigation Report

Lower Fox River and Green Bay, Wisconsin


Prepared for:

Wisconsin Dept. of Natural Resources



Prepared by: The RETEC Group, Inc.
Natural Resource Technology, Inc.

December 2002





Remedial Investigation Report

Lower Fox River and Green Bay, Wisconsin

Prepared by:

**The RETEC Group, Inc.
413 Wacouta Street, Suite 400
St. Paul, Minnesota 55101-1957**

RETEC Project No.: WISCN-14414-345

**Natural Resource Technology, Inc.
23713 W. Paul Road, Unit D
Pewaukee, Wisconsin 53072**

NRT Project No.: 1300

Prepared for:

**Wisconsin Department of Natural Resources
101 S. Webster Street
Madison, Wisconsin 53707-7921**

December 2002

Remedial Investigation Report

Lower Fox River and Green Bay, Wisconsin

Prepared by:

The RETEC Group, Inc.
413 Wacouta Street, Suite 400
St. Paul, Minnesota 55101-1957
RETEC Project No.: WISCN-14414-345

Natural Resource Technology, Inc.
23713 W. Paul Road, Unit D
Pewaukee, Wisconsin 53072
NRT Project No.: 1300

Prepared for:

Wisconsin Department of Natural Resources
101 S. Webster Street
Madison, Wisconsin 53707-7921

Prepared by:



Eric Kovatch, Natural Resource Technology, Inc.

Technically Reviewed by:



Robert Karnauskas, Natural Resource Technology, Inc.



Paul Putzier, The RETEC Group, Inc.

December 2002

Table of Contents

Executive Summary	xix
1 Introduction	1-1
1.1 Project Overview and Objectives	1-1
1.2 Study Area Overview	1-2
1.2.1 Lower Fox River	1-2
1.2.2 Green Bay	1-3
1.3 Study Area River Reaches and Bay Zones	1-6
1.3.1 Lower Fox River Reaches	1-6
1.3.2 Green Bay	1-7
1.3.2.1 Green Bay Zones	1-7
1.3.2.2 Inner and Outer Bays	1-8
1.3.2.3 Lower Green Bay	1-8
1.4 Background	1-8
1.4.1 Site History	1-8
1.4.2 Historical PCB Use and Discharges	1-12
1.4.3 Regulatory Response	1-14
1.4.3.1 Clean Water Act	1-14
1.4.3.2 Wisconsin Pollution Discharge Elimination System	1-14
1.4.3.3 Great Lakes Areas of Concern	1-14
1.5 Application of NRC Findings and Recommendations	1-15
1.6 Section 1 Figures	1-17
2 Database and Investigation Summaries	2-1
2.1 Data Quality Evaluation	2-2
2.2 Sediment Investigations Included in the FRDB	2-3
2.2.1 1989-1990 Fox River Mass Balance Study Data and 1989- 1990 Green Bay Mass Balance Study Data	2-3
2.2.2 1994 Woodward-Clyde Deposit A Sediment Data	2-6
2.2.3 1992/93 BBL Deposit A Sediment Data	2-6
2.2.4 1993 Triad Assessment	2-7
2.2.5 1994 GAS/SAIC Sediment Data	2-8
2.2.6 1995 WDNR Sediment Data	2-8
2.2.7 1996 FRG/BBL Sediment/Tissue Data	2-9
2.2.8 Sediment Remediation Demonstration Projects Data .	2-9
2.2.8.1 Deposit N Demonstration Project	2-11
2.2.8.2 SMU 56/57 Demonstration Project	2-12
2.2.9 1998 FRG/Exponent Data and 1998 FRG/BBL Sediment/Tissue Data	2-15

Table of Contents

2.2.10	1998 RETEC RI/FS Supplemental Data	2-15
2.2.11	Lake Michigan Mass Balance Data	2-17
2.2.12	Fox River Fish Consumption Advisory Data	2-17
2.2.13	USGS National Water Quality Assessment Program (NAWQA) Data	2-17
2.2.14	1997 WDNR Caged Fish Bioaccumulation Study Data	2-17
2.2.15	Minergy Mineralogical Data	2-17
2.3	Ecological Sampling Studies	2-18
2.4	Section 2 Tables	2-19
3	Physical Characteristics	3-1
3.1	Land Use	3-1
3.1.1	Historical Land Use	3-1
3.1.2	Current Land Uses	3-3
3.2	Meteorology	3-5
3.3	Geologic Characteristics	3-6
3.3.1	Regional Geologic Setting	3-7
3.3.1.1	Bedrock Geology	3-7
3.3.1.2	Glacial Geology	3-8
3.3.2	Regional Soils	3-9
3.3.3	Hydrogeology	3-11
3.3.3.1	Regional Hydrogeology	3-11
3.3.3.2	Water Use (1995)	3-15
3.4	Lower Fox River Surface Water Hydrology	3-17
3.4.1	Surface Water Flow Controls	3-17
3.4.1.1	Dams in Wisconsin and on the Lower Fox River	3-17
3.4.1.2	Lower Fox River Dams and Navigational Controls	3-23
3.4.1.3	Neenah-Menasha (Lake Winnebago)	3-24
3.4.2	Lower Fox River Surface Elevation	3-25
3.4.3	Low-Flow and Flood Frequencies	3-26
3.4.4	Measured and Estimated Stream Flow Velocities	3-27
3.4.5	Lower Fox River Bathymetry	3-29
3.4.5.1	LLBdM Reach	3-29
3.4.5.2	Appleton to Little Rapids Reach	3-29
3.4.5.3	Little Rapids to De Pere Reach	3-29
3.4.5.4	De Pere to Green Bay Reach	3-30

Table of Contents

3.5	Green Bay Surface Water Hydrology	3-30
3.5.1	Green Bay Water Level Elevations	3-30
3.5.2	Green Bay Water Circulation, Currents, and Mixing Patterns	3-31
3.5.2.1	Lower Fox River Discharge into Green Bay . . .	3-33
3.5.2.2	Fox River Plume Studies	3-34
3.5.2.3	Inner Bay/Outer Bay Mixing Studies	3-36
3.5.2.4	Green Bay/Lake Michigan Mixing Studies . . .	3-38
3.5.3	Green Bay Bathymetry	3-39
3.5.3.1	Zone 2 Bathymetry	3-41
3.5.3.2	Zone 3 Bathymetry	3-42
3.5.3.3	Zone 4 Bathymetry	3-43
3.5.4	Green Bay Ice Cover	3-45
3.6	Sediment Characteristics	3-46
3.6.1	Sediment Deposition	3-46
3.6.1.1	Lower Fox River Sediment Transport and Deposition	3-46
3.6.1.2	Green Bay Sediment Transport and Deposition	3-49
3.6.1.3	River and Bay Sediment Dredging	3-51
3.6.2	Sediment Grain Size/Lithology	3-52
3.6.3	Estimated Sediment Thickness and Areal Extent . . .	3-53
3.6.3.1	LLBdM Reach	3-55
3.6.3.2	Appleton to Little Rapids Reach	3-55
3.6.3.3	Little Rapids to De Pere Reach	3-55
3.6.3.4	De Pere to Green Bay Reach	3-56
3.6.3.5	Green Bay (Zones 2 through 4)	3-56
3.6.4	Total Organic Carbon	3-56
3.6.5	Other Physical Parameters	3-57
3.7	Section 3 Figures, Tables, and Plates	3-58
4	Ecological Characteristics	4-1
4.1	Overview	4-1
4.1.1	Habitats	4-2
4.1.2	Wildlife Groups	4-3
4.2	Wildlife Habitat	4-5
4.2.1	Open Lands	4-5
4.2.2	Woodlands	4-6
4.2.3	Wetlands	4-8

Table of Contents

4.2.3.1	Wetland Areas and Types	4-8
4.2.3.2	Wetland Losses	4-14
4.2.3.3	Proposed Wetland Restoration Projects	4-18
4.2.4	Riverine Habitat of the Lower Fox River	4-20
4.2.5	Lacustrine Habitat of Green Bay	4-22
4.2.5.1	Overview	4-22
4.2.5.2	Inner Bay Water Quality	4-23
4.2.5.3	Outer Bay Water Quality	4-24
4.3	Benthic Communities	4-25
4.4	Fish	4-27
4.4.1	LLBdM to De Pere Dam Fish Surveys	4-27
4.4.2	De Pere to Green Bay/Duck Creek Fish Surveys	4-28
4.4.3	Green Bay Fishery Observations and Habitat	4-30
4.4.4	Life Histories of Fish Species in the Lower Fox River and Green Bay	4-35
4.4.4.1	Shiners (Minnows)	4-35
4.4.4.2	Gizzard Shad	4-36
4.4.4.3	Rainbow Smelt	4-37
4.4.4.4	Alewife	4-39
4.4.4.5	Yellow Perch	4-41
4.4.4.6	Carp	4-43
4.4.4.7	Walleye	4-45
4.4.4.8	Brown Trout	4-46
4.4.4.9	Sturgeon	4-48
4.5	Birds	4-49
4.5.1	Passerine Birds	4-50
4.5.2	Gulls/Terns	4-51
4.5.3	Diving Birds	4-53
4.5.4	Shorebirds	4-54
4.5.5	Wading Birds	4-55
4.5.6	Waterfowl	4-56
4.5.7	Raptors	4-57
4.6	Mammals	4-59
4.6.1	Mink	4-60
4.6.1.1	Mink Habitat	4-60
4.6.1.2	Domestic Mink Production in Wisconsin	4-61
4.6.1.3	Wild Mink in the Study Area	4-62
4.6.2	Otter	4-62

Table of Contents

4.7	Endangered and Threatened Species	4-63
4.8	Section 4 Figures and Tables	4-64
5 Nature and Extent of Detected Chemicals		5-1
5.1	Detected Compound Sources	5-1
5.1.1	Point Sources	5-1
5.1.1.1	Industrial/Municipal Discharges	5-1
5.1.1.2	Landfills	5-5
5.1.1.3	Spills	5-6
5.1.2	Non-Point Sources	5-7
5.1.2.1	River and Bay Sediments	5-8
5.1.2.2	Stormwater Runoff	5-9
5.1.2.3	Atmospheric Deposition and Volatilization ...	5-11
5.2	Summary of Detected Chemicals	5-12
5.2.1	Overview	5-12
5.2.2	PCBs	5-14
5.2.3	Dioxin/Furan	5-18
5.2.4	Pesticides	5-19
5.2.5	Inorganic Compounds	5-19
5.2.6	TCLP Results	5-22
5.2.7	Semi-Volatile Organic Compounds (SVOCs)	5-22
5.3	Lake Winnebago (Background) Results	5-23
5.4	Chemical Distribution in Sediments	5-24
5.4.1	Overview	5-24
5.4.2	PCB Distribution	5-25
5.4.2.1	Bed Maps and Sediment Data Interpolation Methods	5-25
5.4.2.2	Lower Fox River and Green Bay PCB Results ..	5-31
5.4.2.3	LLBdM Reach PCB Results	5-32
5.4.2.4	Appleton to Little Rapids Reach PCB Results ..	5-36
5.4.2.5	Little Rapids to De Pere Reach PCB Results ..	5-40
5.4.2.6	De Pere to Green Bay Reach PCB Results	5-42
5.4.2.7	Green Bay Zone 2 PCB Results	5-46
5.4.2.8	Green Bay Zone 3 PCB Results	5-48
5.4.2.9	Green Bay Zone 4 PCB Results	5-50
5.4.2.10	General PCB Homolog Distribution	5-52
5.4.3	Dioxin/Furans	5-55
5.4.4	Pesticides	5-56

Table of Contents

5.4.4.1	LLBdM Reach Results	5-56
5.4.4.2	Appleton to Little Rapids Reach Results	5-57
5.4.4.3	Little Rapids to De Pere Reach Results	5-57
5.4.4.4	De Pere to Green Bay Reach Results	5-57
5.4.5	Inorganic Compounds	5-58
5.4.5.1	Mercury	5-58
5.4.5.2	Lead	5-59
5.4.5.3	Arsenic	5-61
5.4.6	Other Organic Compounds	5-62
5.4.6.1	LLBdM Reach SVOC Results	5-62
5.4.6.2	Appleton to Little Rapids Reach SVOC Results	5-63
5.4.6.3	Little Rapids to De Pere Reach SVOC Results	5-63
5.4.6.4	De Pere to Green Bay Reach SVOC Results	5-64
5.4.6.5	Green Bay SVOC Results	5-64
5.4.7	Other Inorganic Compounds	5-64
5.4.7.1	Cadmium/Chromium	5-64
5.4.7.2	Ammonia	5-65
5.5	Surface Water Sampling Results	5-65
5.5.1	Overview	5-65
5.5.2	PCB Distribution	5-66
5.5.2.1	Distribution in the Lower Fox River	5-67
5.5.2.2	PCB Distribution in Green Bay	5-69
5.5.2.3	PCB Distribution in Lake Michigan	5-69
5.5.3	Mercury Distribution	5-70
5.5.4	Pesticide Distribution	5-70
5.6	Chemical Loading to Green Bay	5-71
5.6.1	PCB Loading to Green Bay	5-71
5.6.2	Mercury and DDT Loading to Green Bay	5-71
5.7	Summary of PCBs in Biota	5-72
5.8	Time Trends of Contaminants in Sediment and Fish	5-72
5.8.1	Sediment Methods	5-72
5.8.2	Fish Methods	5-74
5.8.3	Results	5-75
5.8.3.1	Little Lake Butte des Morts	5-76
5.8.3.2	Appleton to Little Rapids	5-78
5.8.3.3	Little Rapids to De Pere	5-78
5.8.3.4	De Pere to Green Bay (Zone 1)	5-79
5.8.3.5	Green Bay Zone 2	5-80

Table of Contents

5.8.4	Conclusion	5-80
5.9	Section 5 Figures, Tables, and Plates	5-82
6	Chemical Transport and Fate	6-1
6.1	Introduction	6-1
6.2	Transport and Fate Processes	6-1
6.2.1	Chemical Transport Interactions	6-1
6.2.2	Physical Transport	6-3
6.2.3	Biological Interactions	6-4
6.3	Compounds of Potential Concern	6-5
6.3.1	Organic Constituents	6-5
6.3.1.1	PCBs	6-5
6.3.1.2	Dioxins and Furans	6-6
6.3.1.3	DDT	6-7
6.3.1.4	Dieldrin	6-8
6.3.2	Inorganic Constituents	6-9
6.3.2.1	Mercury	6-9
6.3.2.2	Lead	6-10
6.3.2.3	Arsenic	6-11
6.4	Lower Fox River/Green Bay Modeling	6-12
6.4.1	GBTOXe Model	6-12
6.4.2	GBFood Model	6-12
6.4.3	Fox River Food (FRFood) Model	6-13
6.4.4	Whole Lower Fox River Model	6-14
6.5	Section 6 Tables	6-15
7	Summary of Findings	7-1
7.1	Introduction	7-1
7.2	Physical and Ecological Characteristics	7-2
7.3	Nature and Extent of Sediment Impacts	7-4
7.3.1	Overview	7-4
7.3.2	Lower Fox River PCB Impacts	7-6
7.3.2.1	Overview	7-6
7.3.2.2	Little Lake Butte des Morts Reach	7-6
7.3.2.3	Appleton to Little Rapids Reach	7-7
7.3.2.4	Little Rapids to De Pere Reach	7-8
7.3.2.5	De Pere to Green Bay Reach	7-8
7.3.3	Green Bay PCB Impacts	7-9

Table of Contents

7.3.3.1	Overview	7-9
7.3.3.2	Green Bay Zone 2	7-10
7.3.3.3	Green Bay Zone 3	7-11
7.3.3.4	Green Bay Zone 4	7-12
7.3.4	Other Chemical Compounds	7-12
7.4	Chemical Transport and Fate	7-15
7.5	Investigative Assumptions/Uncertainties	7-16
8	References	8-1

Table of Contents

Appendix A	Data Management Summary Report (EcoChem, 2000) Addendum 1 to the Data Management Summary Report (EcoChem, 2002)
Appendix B	Time Trends Analysis (Mountain-Whisper-Light, 2001)
Appendix C	A Windrose diagram, developed from the National Oceanic and Atmospheric Administration
Appendix D	Stratigraphic cross-sections and other pertinent information concerning the regional geology of the area (Krohelski and Brown, 1986).
Appendix E	Bathymetry information available from the NOAA recreational charts for Lake Winnebago and the Lower Fox River (NOAA, 1992)
Appendix F	The USGS hydrographs for two storm events in November 1998
Appendix G	Physical parameters tables
Appendix H	PCB Congeners and Homolog Group List (ATSDR, 1997a)
Appendix I	PCB congener results for each reach and zone

List of Figures

- Figure 1-1 Lower Fox River Study Area
- Figure 1-2 Green Bay Study Area
- Figure 1-3 Little Lake Butte des Morts Reach
- Figure 1-4 Appleton to Little Rapids Reach
- Figure 1-5 Little Rapids to De Pere Reach
- Figure 1-6 De Pere to Green Bay Reach
- Figure 3-1 Lower Fox River Elevation Profile
- Figure 3-2 Green Bay Monthly Mean Surface Circulation - August 1989
- Figure 3-3 Green Bay Monthly Bottom Surface Circulation - August 1989
- Figure 3-4 Green Bay Zone 2 Bathymetry
- Figure 3-5 Green Bay Zone 3 Bathymetry
- Figure 3-6 Green Bay Zone 4 Bathymetry
- Figure 3-7 Estimated Annual Sediment Transport Rates and Stream Flow Velocities
- Figure 4-1 Lower Fox River Wetland, Habitat, and Animal Distribution: Little Lake Butte des Morts Reach
- Figure 4-2 Lower Fox River Wetland, Habitat, and Animal Distribution: Appleton to Little Rapids Reach
- Figure 4-3 Lower Fox River Wetland, Habitat, and Animal Distribution: Little Rapids to De Pere Reach
- Figure 4-4 Lower Fox River Wetland, Habitat, and Animal Distribution: De Pere to Green Bay Reach
- Figure 4-5 Wetland Distribution: Green Bay Zones 2 and 3
- Figure 4-6 Wetland Distribution: Green Bay Zone 4
- Figure 4-7 Wetland Losses in Green Bay: Duck Creek, Cat Island Chain, and Long Tail Point
- Figure 4-8 Green Bay Spawning Areas by Fish Types: Salmon/Trout and Benthic Fish
- Figure 4-9 Green Bay Spawning Areas by Fish Types: Pelagic and Game Fish
- Figure 4-10 Green Bay Spawning Areas by Fish Species: Walleye, Yellow Perch, and Sturgeon
- Figure 4-11 Green Bay Spawning Areas by Fish Species: Carp and Alewife
- Figure 4-12 Green Bay Spawning Areas by Fish Species: Emerald Shiners and Gizzard Shad
- Figure 4-13 Distribution of Birds in Green Bay: Select Species and Groups
- Figure 4-14 Lower Fox River Mink Habitat Suitability: Little Lake Butte des Morts Reach
- Figure 4-15 Lower Fox River Mink Habitat Suitability: Appleton to Little Rapids Reach
- Figure 4-16 Lower Fox River Mink Habitat Suitability: Little Rapids to De Pere Reach
- Figure 4-17 Lower Fox River Mink Habitat Suitability: De Pere to Green Bay Reach
- Figure 4-18 Green Bay Mink Habitat Suitability: Zone 2
- Figure 4-19 Green Bay Mink Habitat Suitability: Zone 3

List of Figures

- Figure 5-1 PCB Sampling Frequency Distribution in Lower Fox River and Green Bay Sediments
- Figure 5-2 Summary of Total PCB Concentrations in Lower Fox River and Green Bay Sediments
- Figure 5-3 PCB Mass Distribution in Sediments for Each River Reach and Bay Zone
- Figure 5-4 PCB Mass by Concentration Ranges in Lower Fox River Sediments
- Figure 5-5 Contaminated Sediment Volume by Concentration Ranges in the Lower Fox River
- Figure 5-6 PCB Mass by Concentration Ranges in Green Bay Sediments
- Figure 5-7 Contaminated Sediment Volume by Concentration Ranges in Green Bay
- Figure 5-8 PCB Mass Distribution in Lower Fox River and Green Bay Sediments
- Figure 5-9 Ratios of PCB Mass/Sediment Volume in the Lower Fox River and Green Bay Sediment
- Figure 5-10 Distribution of PCB Mass/Sediment Volume Ratios in Sediments with More Than 50 $\mu\text{g/kg}$ PCBs
- Figure 5-11 Aroclors 1242/1254/1260 PCB Homolog Plots
- Figure 5-12 PCB Homolog Distribution in Lower Fox River and Green Bay Sediments
- Figure 5-13 1994/95 Total PCB & Mercury Concentrations in Lower Fox River Water
- Figure 5-14 1998 Total PCB Concentrations in Lower Fox River Water
- Figure 5-15 PCB Homolog Distribution in Water: De Pere Dam Through Green Bay
- Figure 5-16 Lower Fox River and Green Bay System - Estimated PCB Mass and Major PCB Flux Pathways
- Figure 5-17 Time Trends of PCBs in Sediments for Depths from 0 to 10 cm and from 10 to 30 cm
- Figure 5-18 Time Trends of PCBs in Sediments for Depths from 30 to 50 cm and from 50 to 100 cm
- Figure 5-19 Time Trends of PCBs in Sediments for Depths over 100 cm

List of Tables

Table 2-1	Fox River Database Studies and Data Classification
Table 2-2	Lower Fox River - Supplemental Data Collection Sampling List
Table 3-1	Land Use Classification for Counties Bordering Green Bay
Table 3-2	Temperature and Precipitation Data for the City of Green Bay, Wisconsin
Table 3-3	Temperature and Precipitation Data for the City of Appleton, Wisconsin
Table 3-4	Temperature and Precipitation Data for the City of Marinette, Wisconsin
Table 3-5	Temperature and Precipitation Data for the City of Sturgeon Bay, Wisconsin
Table 3-6	Temperature and Precipitation Data for the City of Fayette, Michigan
Table 3-7	Water Use in the Lower Fox River/Green Bay Watersheds (1995)
Table 3-8	Lower Fox River Dams
Table 3-9	Lower Fox River - U.S. Army Corps of Engineers - Dam Stability and Inspection Information
Table 3-10	Lower Fox River Gradient and Lock/Dam Information
Table 3-11	Lower Fox River Discharge Results - Rapide Croche Gauging Station
Table 3-12	Lower Fox River Stream Velocity Estimates
Table 3-13	Fox River Mouth Gauging Station Results (1989-1999)
Table 3-14	Lower Fox River Total Suspended Solid (TSS) Loads
Table 3-15	USACE Navigation Channel Dredging Records (1957-1999)
Table 4-1	Major Green Bay Wetland Areas/Complexes
Table 4-2	Lower Fox River Habitats
Table 4-3	Lower Fox River Shoreline and Substrate Types
Table 4-4	Lower Fox River Fish Species Composition
Table 4-5	Lower Fox River Fish Populations in the De Pere to Green Bay Reach
Table 4-6	Green Bay Fish Species
Table 4-7	Lower Fox River and Green Bay Bird Species
Table 5-1	Lower Fox River and Green Bay - Sediment Sampling Results: Summary of Detected Compounds
Table 5-2	Lower Fox River and Green Bay - Distribution of Total PCBs in Sediment
Table 5-3	Lower Fox River - Dioxin/Furan (2,3,7,8-TCDD/F) Results
Table 5-4	Lower Fox River and Green Bay - Pesticide Results
Table 5-5	Lower Fox River and Green Bay - Mercury, Lead, and Arsenic Results
Table 5-6	Lower Fox River and Green Bay - Other RCRA Metals, Copper, Nickel, and Zinc
Table 5-7	Lower Fox River and Green Bay - Miscellaneous Inorganic Compounds
Table 5-8	Lower Fox River and Green Bay - Ammonia Results
Table 5-9	Lower Fox River - Toxicity Characteristic Leaching Procedure (TCLP) Results
Table 5-10	Lower Fox River and Green Bay - Semi-Volatile Organic Compound Results (PAHs)
Table 5-11	Lower Fox River and Green Bay - Miscellaneous SVOC Results

List of Tables

Table 5-12	Lake Winnebago Background Sediment Results
Table 5-13	Lower Fox River - PCB Mass and Sediment Volume by Concentration Range
Table 5-14	Lower Fox River - PCB Mass and Sediment Volume by Deposit/SMU Layer
Table 5-15	Green Bay - PCB Mass and Sediment Volume by Concentration Range and Layer
Table 5-16	Lower Fox River and Green Bay - Water Sampling Results: Summary of Detected Compounds
Table 5-17	Lower Fox River - Total PCB Results in Water
Table 5-18	Green Bay - Total PCB Results in Water
Table 5-19	Lower Fox River and Green Bay - Mercury and DDT (DDD/DDE) Water Sampling Results
Table 5-20	PCB Transport within the Lower Fox River and Green Bay System
Table 5-21	Distribution of Resident Tissue Samples over Time in the Lower Fox River - Total PCBs Only
Table 5-22	Distribution of Resident Tissue Samples over Time in Green Bay - Total PCBs Only
Table 5-23	Results of Sediment Time Trends Analysis for the Lower Fox River
Table 5-24	Mass-Weighted Combined Time Trend for 0 to 10 cm Depth by Reach
Table 5-25	Results of Fish Time Trend Analysis on the Lower Fox River
Table 6-1	Lower Fox River - Fate and Transport Chemical Factors

List of Plates

Plate 3-1	Sample Locations and Interpolated Thickness of Sediment with PCBs: Little Lake Butte des Morts Reach
Plate 3-2	Sample Locations and Interpolated Thickness of Sediment with PCBs: Appleton to Little Rapids Reach
Plate 3-3	Sample Locations and Interpolated Thickness of Sediment with PCBs: Little Rapids to De Pere Reach
Plate 3-4	Sample Locations and Interpolated Thickness of Sediment with PCBs: De Pere to Green Bay Reach
Plate 3-5	Sample Locations and Interpolated Thickness of Sediment with PCBs: Green Bay
Plate 5-1	Interpolated PCB Distribution in Sediments: Little Lake Butte des Morts Reach
Plate 5-2	Interpolated PCB Distribution in Sediments: Appleton to Little Rapids Reach
Plate 5-3	Interpolated PCB Distribution in Sediments: Little Rapids to De Pere Reach
Plate 5-4	Interpolated PCB Distribution in Sediments: De Pere to Green Bay Reach
Plate 5-5	Interpolated PCB Distribution in Sediments: Green Bay

List of Acronyms

AOC	Areas of Concern
ATSDR	Agency for Toxic Substances and Disease Registry
AVM	acoustical velocity meter
BCF	bioconcentration factor
BBL	Blasland, Bouck, & Lee
BEHP	bis2-ethylhexylphthalate
°C	degrees Celsius
CDF	confined disposal facility
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
cfs	cubic feet per second
cm	centimeter(s)
COPC	chemical of potential concern
CWA	Clean Water Act
DDD	p,p'-Dichlorodiphenyldichloroethane
DDE	p,p'-Dichlorodiphenyldichloroethylene
DDT	p,p'-Dichlorodiphenyltrichloroethane
DO	dissolved oxygen
ECWRPC	East Central Wisconsin Regional Planning Commission
EIS	Final Environmental Impact Statement
EPA	United States Environmental Protection Agency, Region 5
Exponent	Exponent Environmental Group
°F	degrees Fahrenheit
F&VD	Foth & Van Dyke
FRRAT	Fox River Remediation Advisory Team
FRDB	Fox River Database
FRG	Fox River Group, which is comprised of the following seven companies listed alphabetically: Appleton Papers, Inc., Fort James Corporation; P.H. Glatfelter Company; NCR Corporation; Riverside Paper Corporation; U.S. Paper Mills Corporation; and Wisconsin Tissue Mills, Inc.
FS	Feasibility Study
ft	foot (feet)
ft ²	square feet
ft/s	foot per second
g	gram(s)
g/year	grams per year
GAS	Graef, Anhalt, Schloemer, and Associates, Inc.
GBMBS	Green Bay Mass Balance Study
GBMSD	Green Bay Metropolitan Sewerage District
GBTOXe	Green Bay Toxics Model

List of Acronyms

GIS	Geographic Information System
GLC	Great Lakes Commission
GLNPO	USEPA Great Lakes National Program Office
GLWQA	Great Lakes Water Quality Agreement
gpm	gallons per minute
HDPE	high-density polyethylene
IGLD 1985 Canada	International Great Lakes Datum, zero elevation at Rimouski, Quebec,
IJC	International Joint Commission
in	inch(es)
IPS	Integrated Papers Services
kg	kilogram
km	kilometer(s)
km ²	square kilometers
km ³	cubic kilometers
Koc	log water/organic carbon partition coefficient
Kow	log octanol/water partition coefficient
LLBdM	Little Lake Butte des Morts
LTA	long-term average
LUST	leaking underground storage tank
m	meter(s)
m ²	square meters
m/s	meters per second
m ³	cubic meters
m ³ /s	cubic meters per second
mg/kg	milligrams per kilogram
mg/L	milligrams per liter
MGD	million gallons per day
mi	mile(s)
mi ²	square miles
mi ³	cubic miles
MNFI	Michigan Natural Features Inventory
MSA	Metropolitan Statistical Area
MT	metric tonnes
NWR	national wildlife refuge
NAWQA	USGS National Water Quality Assessment Program
NCP	National Contingency Plan
NCR	National Cash Register
ng/kg	nanograms per kilogram

List of Acronyms

ng/L	nanograms per liter
NOAA	National Oceanic and Atmospheric Administration
NPDES	National Pollution Discharge Elimination System
NPL (Superfund)	National Priority List
NRC	National Research Council
NURE	National Uranium Resources Evaluation project
NWR	National Wildlife Refuge
OZ	ounce
PAH	polynuclear aromatic hydrocarbon
PCB	Polychlorinated Biphenyl
PCP	Pentachlorophenol
pg/m ³	picograms per cubic meter
ppb	parts per billion ($\mu\text{g/kg}$ or $\mu\text{g/L}$)
ppm	parts per million (mg/kg or mg/L)
ppt	part per trillion (ng/kg or ng/L)
Project Team	The Fox River Project Team
PRP	Principal Responsibility Parties
PSC	Public Service Commission
QAPP	Quality Assurance Project Plan
QA/QC	Quality Assurance/Quality Control
RA	Risk Assessment
RAP	Remedial Action Plan
RCRA	Resource Conservation and Recovery Act
RI	Remedial Investigation
SAIC	Science Application International Corporation
SAV	submerged aquatic vegetation
SCS	Soil Conservation Service
SEF	Sediment enrichment factor
SLRA	Screening Level Risk Assessment
SMU	Sediment Management Unit
SQG	sediment quality guideline
SRD	Sediment Remediation Demonstration
SVOC	semi-volatile organic compound
SWA	State Wildlife Area
SWE	Snow-Water Equivalent
TCLP	Toxicity Characteristic Leaching Procedure
TOC	total organic carbon
TRI	EPA Toxic Release Inventory database
TSCA	Toxic Substance Control Act

List of Acronyms

TSS	total suspended solids
$\mu\text{g/kg}$	microgram per kilogram
UP	Michigan's Upper Peninsula
USACE	U.S. Army Corps of Engineers
USFWS	U.S. Fish and Wildlife Service
USGS	United States Geological Survey
UST	underground storage tank
UWSGI	University of Wisconsin Sea Grant Institute
VOC	volatile organic compound
W.A.	Wildlife Area
W.A.C.	Wisconsin Administrative Code
WCC	Woodward-Clyde Consultants (formerly EWI Engineering Associates)
WDNR	Wisconsin Department of Natural Resources
WGNHS	Wisconsin Geological and Natural History Survey
WPDES	Wisconsin Pollution Discharge Elimination System
WPSC	Wisconsin Public Service Corporation
WSCO	Wisconsin State Climatology Office
WTM 27	Wisconsin Trans-Mercator Projection, 1927
WWTP	Wastewater Treatment Plant
WY	Water Year
yd^3	cubic yard
YOY	Young-of-year fish

EXECUTIVE SUMMARY

The Remedial Investigation (RI) report summarizes the physical, chemical, and biological characteristics of the Lower Fox River and Green Bay. The purpose of the RI report is to compile and evaluate these data to support development of the Baseline Human Health and Ecological Risk Assessment (RA) and Feasibility Study (FS). The RA identifies the risks posed to human health and the environment by compounds of concern. The FS develops and evaluates a range of remedial alternatives to support the selection of a remedy that will eliminate, reduce and/or control these risks. This RI/FS report is consistent with the findings of the National Academy of Science's National Research Council Report entitled *A Risk Management Strategy for PCB Contaminated Sediments*. (NRC, 2001).

The RI study area includes the Lower Fox River extending 63 km (39 mi) from Lake Winnebago to Green Bay as well as the entire 4,150 km² (1,600 mi²) of the bay. Green Bay is 190 km (119 mi) in length and averages 37 km (23 mi) in width. The Lower Fox River was subdivided into four river reaches. Green Bay is subdivided into zones 2, 3, and 4 (Figure 1). The Green Bay Area of Concern, as designated by the International Joint Commission, is defined as the De Pere to Green Bay Reach and much of Green Bay Zone 2.

The RI evaluated data from numerous investigations conducted within the study area since 1971, which comprise the Fox River Database (FRDB). Sediment, water, and biological samples in the FRDB include analyses for over 200

chemical parameters. Based on these analyses, a Screening Level Risk Assessment identified polychlorinated biphenyls (PCBs), dieldrin, DDT, dioxins/furans, mercury, lead, and arsenic as the compounds present in the study area that represent potential risks to human health and the environment. However, PCBs are the primary compounds of concern.




Figure 1. Remedial Investigation Study Areas

Site History and PCB Discharges

In the early 1950s, carbonless copy paper was developed through a process that applied an emulsion containing PCB on paper in a manner that would create document copies. Lower Fox River valley paper mills manufactured and recycled this carbonless paper between 1954 and 1971. About 45 million pounds of PCB were used in the Fox Valley during this time period.

PCBs were released to the environment through manufacturing waste waters and from the de-inking/recycling of waste carbonless copy paper. The Wisconsin Department of Natural Resources (WDNR) estimates the amount of PCB that was discharged to the Lower Fox River from these activities is 313,600 kg (691,370

pounds), with a range from 126,450 kg to 399,450 kg (278,775 to 880,640 pounds). WDNR believes that five facilities contributed over 99 percent of the total PCBs discharged to the Lower Fox River by the end of 1971.

In the late 1970s, commercial production of PCBs in the United States was prohibited due to concerns for human health and the environment. At the present time, some minor unavoidable point source discharges along with atmospheric deposition of PCB continue, but are small compared to the PCB mass present in the river and bay sediments.

Prior to implementation of the federal Clean Water Act in 1972, rough fish were the main species that could live in the Lower Fox River. With implementation of the Clean Water Act and more stringent control over wastewater discharges, water quality in the river improved and game fish began to return to the river. PCBs were detected in trout from Green Bay as early as 1971. Due to continued elevated PCB levels, WDNR issued advisories for public consumption of fish (1976) and waterfowl (1983) derived from Green Bay and the Lower Fox River. The state of Michigan also issued consumption advisories for Green Bay fish in 1977.

PCB Distribution and Sediment Volumes

Considering sediments containing more than 50 $\mu\text{g/kg}$ PCB, about 28,600 kg (63,050 pounds) of PCBs are contained within about 9 million m^3 (11.8 million yd^3) of sediment in the Lower Fox River. In Green Bay, approximately 68,200 kg (150,300 pounds) of PCBs are dispersed in about 465 million m^3 (610 million

yd^3) of sediment. The distribution of PCB mass, sediment volume and sediment areal extent are shown on Figure 2. Also shown on Figure 2 is the ratio of PCB mass to sediment volume. The reaches upstream of the De Pere dam are combined on Figure 2 because of their relatively small PCB mass, sediment volume and areal extent.

Much of the PCB discharged into the Lower Fox River has already been transported downstream and is now concentrated in sediments within specific areas:

- The De Pere to Green Bay Reach contains almost 26,000 kg of PCB, which represents about 91 percent of the mass remaining in the river. This reach contains just under 27 percent of the total PCB mass in the system and is concentrated within a relatively small area comprising just over one percent of the total sediment volume. This reach also exhibits the highest mass of PCB per volume of sediment.
- Approximately 70 percent of the total PCB mass in the system has migrated from the river into Green Bay.
- The PCB mass in Green Bay is dispersed over an extraordinarily large area and in an extremely large sediment volume. Almost half of the total PCB mass in Green Bay is found in Zone 2.

Sediment and PCB Transport

Particle size and cohesion along with river/bay conditions, especially current speeds, control the deposition, resuspension, and transportation of sediments (and the PCBs absorbed to them). In the Lower Fox River, sediments have accumulated in 35 separate deposits above




Figure 2. PCB Mass & Sediment Volume/Area Distribution by river reach and bay zone

the De Pere dam. Below the De Pere dam and in Green Bay, where current speeds tend to be lower, sediments cover large areas of the river and bay bottom, except in areas where the sediments are dredged to maintain ship navigation. The highest PCB concentrations have also been observed in the LLBdM and De Pere to Green Bay reaches, in the vicinity of historic discharge points.

The average river discharge was about 122 m³/s (4,300 cfs) between 1989 and 1998. Due to storm events and spring snowmelt, the river discharge exceeds 272 m³/s (9,600 cfs) more than 10 percent of the time. These faster currents have the capability to resuspend and transport larger particle sizes and greater volumes of sediment and, therefore, a greater mass of PCB. Field measurements and computer modeling results suggest that

these less-frequent, high-discharge events transport much of the PCB mass in the river over the De Pere dam and into Green Bay. In addition to sediment transport, PCB migrates due to dissolution in water and adsorption onto algae and other organic matter. The PCB mass transported from reach to reach increases along the river. Based on sampling data collected as part of the Green Bay Mass Balance study in 1989-90, about 280 kg (610 pounds) of PCB were transported to Green Bay during the study period. Based on work done in 1994-95 as part of the Lake Michigan Mass Balance, it was estimated that 220 kg (485 pounds) of PCB moved from the river into the bay. PCB loads to the bay vary as the river flow varies. This mass represents up to 1 percent of the PCB mass in the river.

Sediment discharged from the Lower Fox River is directed toward the east shore of Green Bay by counterclockwise currents. This sediment-rich water can extend between 20 km to 40 km (12 mi to 24 mi) along the east shore. Fluctuating water levels, wave action and reverses in stream flow in this area facilitate sediment transport and mixing. Consequently, large volumes of sediment containing PCB are present along the southern and eastern portions of Green Bay. At least 68,200 kg (150,300 pounds) of PCBs already reside in the bay. Over 95 percent of the PCB that occurs in Green Bay is derived from the Lower Fox River.

This transport of PCB also extends into Lake Michigan. During 1989/90, it was estimated as part of the Green Bay Mass Balance Study that about 122 kg (270 pounds) of PCBs were transported from Green Bay to Lake Michigan. Other mass transport pathways (such as volatilization) also exist.

Ecological Samples and Characteristics

Exposure of biota to sediments and water containing PCB fosters uptake of PCBs into the food chain. Wetlands, submerged aquatic vegetation, and islands along the Lower Fox River and Green Bay offer nesting/spawning, feeding, and refuge opportunities for fish, birds, and animals. Other lacustrine, riverine, and estuary features also provide habitat for regional wildlife. In addition to birds and fish, the FRDB contains information on PCBs in deer, otter, mink, and various insects and invertebrates. The RA evaluates PCB uptake and accumulation in selected species and the associated human health and

environmental risks. Areas with higher PCB concentrations tend to pose a greater risk of exposure.

Effects of Time

The FRDB includes sediment and water results from over a 10 year period while tissue samples were collected between 1971 and 1999. During the 1970s, after PCB discharges into the river ceased, PCB concentrations in fish tissue showed significantly declining concentrations. However, since the mid-1980s, changes in the rate of PCB decline in fish tissue have been observed. Changes in PCB levels in fish tissue have either slowed, remained constant, or in some cases actually increased.

PCB concentration trends in the upper 10 cm (4 in) of sediment are inconsistent, but generally appear to be decreasing over time as more PCB is transported downstream. Soil eroded from the watershed mixes with and may further dilute PCB concentrations in the sediments.

Further Information

The selection of remedies for the Lower Fox River and Green Bay will consider the information within the RI, RA and FS, as well as input by the public and interested parties. For further information, please contact:

Mr. Edward K. Lynch, P.E.
WI Department of Natural Resources
101 S. Webster Street
Box 7921, Madison, WI 53703
(608/266-3084)
or visit the WDNR website at
<http://www.dnr.state.wi.us/org/water/wm/lowerfox>